

## **Guidance on the management of transplant recipients diagnosed with or suspected of having COVID19**

Updated 24<sup>th</sup> April 2020

### **AIMS**

To provide guidance on the management of transplanted patients diagnosed with or suspected of having COVID 19, based on the limited available evidence.

### **OVERVIEW OF GUIDANCE**

This is a consensus opinion of a group of transplant professionals. The guidance is based on the sparse information available on transplant recipients in the literature and should be used in conjunction with local or national guidance. We understand that individual patients may need a bespoke plan but this is a general guideline which may help others when managing transplant recipients with, or suspected of having, COVID 19.

All unwell transplant recipients should be discussed with their usual transplant unit.

This guidance does not cover cardiothoracic or small bowel transplant recipients.

Caution should be exercised when reducing immunosuppression in paediatric recipients and all cases must be discussed with their paediatric transplant centres first.

At the time of writing, there are no randomised controlled trials of additional therapy that show any benefit above standard supportive care.

The guidance is correct as of April 16th 2020 and may be updated further.

### **EXECUTIVE SUMMARY**

- Exclude all other causes of fever and symptoms
- Discontinue antiproliferative agents (Aza/MMF) and restart after full recovery
- Calcineurin inhibitors
  - Review and minimise in early disease
  - Reduce or discontinue in severe or progressive disease
- High dose steroid
  - can be counterproductive in early disease
  - may be considered in progressive pulmonary disease/ARDS
- Antivirals, chloroquine/hydroxychloroquine or biologics should be used in line with local protocols or research studies
- Monitor inpatients closely for rapid deterioration
- Maintain circulating volume but avoid excessive fluid administration
- Early thromboprophylaxis is recommended, unless contraindications
- All cases in transplanted or waiting list patients should be reported to NHS BT

## GENERAL PRINCIPLES

Staff evaluating transplant patients with fever or cough should:

- follow national or local guidelines on the use of personal protective equipment
- exclude other causes for symptoms (Eg CMV, pneumocystis, community or hospital acquired pneumonia, influenza, urinary sepsis, lymphoma and fluid overload amongst other diagnoses)
- consider atypical presentations of COVID (eg loin pain in patients with lower lobe infection) and have a low threshold for considering COVID
- Naso pharyngeal swab for PCR analysis is recommended in all potential cases
- Staff must only perform swabbing if they are trained in acquisition and have appropriate PPE. [See NHS BT guidance](#)
- Nasopharyngeal swab is specifically indicated if:
  - An alternative treatable cause is considered (eg Influenza)
  - It is required for diagnostic certainty – in admitted or deteriorating patients
  - It is required for purposes of isolation
- A negative swab result requires repeat if clinical suspicion is high
- All proven or suspected cases should be reported to NHS BT
  - <https://www.odt.nhs.uk/deceased-donation/covid-19-advice-for-clinicians/>

## USEFUL LINKS

The guidance is not intended to be comprehensive and UK national guidance links are below:

<https://www.gov.uk/government/publications/wuhan-novel-coronavirus-initial-investigation-of-possible-cases>

NHS England guidance to clinicians can be found here:

<https://www.england.nhs.uk/coronavirus/>

Detailed clinical guidance from the WHO can be found here:

[https://www.who.int/publications-detail/clinical-management-of-severe-acute-respiratory-infection-when-novel-coronavirus-\(ncov\)-infection-is-suspected](https://www.who.int/publications-detail/clinical-management-of-severe-acute-respiratory-infection-when-novel-coronavirus-(ncov)-infection-is-suspected)

## **CLINICAL GUIDELINES FOR MANAGING PATIENTS WITH OR SUSPECTED OF HAVING COVID 19**

**See appendix 1 for flow chart**

### **1: Patients who do not require hospital admission**

The majority of patients will have mild symptoms and do not require admission. Each patient should be considered individually regarding the risk of immunosuppression dose reduction. Particular care should be exercised in paediatric recipients who may have a milder illness.

Experience from international centres suggests the discontinuation of antiproliferatives, such as mycophenolate and azathioprine, in line with clearance of other viral pathogens.

Experimental evidence suggests that coronavirus may require intact immunophilin pathways with a role for tacrolimus and cyclosporin to inhibit the growth of human coronaviruses<sup>1,2</sup>. This data would suggest that calcineurin inhibitors may be the agent of choice to continue but their potential benefit has to be balanced against their immunosuppressive effect. Levels should be reviewed and minimised in the face of active infection.

High dose steroids have previously been suggested as therapy for coronavirus lung injury but are associated with prolonged viral shedding<sup>3</sup> and may have deleterious effect in experimental models<sup>4</sup>. Early use of hydrocortisone in SARS CoV1 was associated with higher viral loads<sup>5</sup> and more recent analysis does not support their use, unless required for other indications<sup>6</sup>.

Patients in the early phase may have false negative naso pharyngeal swab results (~30%) and repeat nasal swab may be required if first result is negative and clinical suspicion is high, particularly if a result is required for isolation protocols. It is important to review the naso pharyngeal swab result in conjunction with clinical suspicion and radiological changes.

#### **Recommendation:**

- Stop antiproliferative agents (MMF/azathioprine)
- Review total burden of immunosuppression and consider reduction of CNI
- High or increased dose steroid is NOT recommended at this stage
- Patients should self isolate in line with national guidance
- Closely monitor patients remotely for change in symptoms
- Consider restarting immunosuppression 14 days after onset of symptoms if symptom free in absence of anti-pyretics for minimum of 3 days
- Consider early monitoring of graft function when safe to do so and risk of transmission to others is low
- Due to prolonged viral shedding in immunosuppressed patients and the risk of disease transmission, post COVID transplant recipients must be reviewed in an area separate from non-affected individuals.

## 2: Patients who are unwell and admitted to hospital

Initial reports from the Chinese Centre for Disease Control and prevention suggest that 81% of non-immunosuppressed patients have a mild disease with 14% having severe disease and 5% requiring intensive care support<sup>7</sup>. The overall mortality was 2.3%, rising to 49% in patients requiring critical care. There is insufficient data to draw a firm conclusion on the course of transplanted patients, although one study from Spain<sup>14</sup> in a variety of hospitalised transplant recipients recorded a 27.8% case fatality and very early UK data in renal transplant recipients suggests a mortality of 12-16%<sup>15</sup>. In a further study of 138 hospitalised patients from China, the median onset of dyspnoea from first symptoms was 5 days, with admission on day 7 and onset of ARDS at day 8<sup>8</sup>. The scenario of rapid deterioration and requirement for ventilation one week after onset of symptoms is an increasingly recognised course for COVID.

Initial reports from China suggested Acute Kidney Injury rates around 3%<sup>8,10</sup> with higher mortality rates in patients with renal impairment<sup>10</sup>. [See summary.](#) Initial UK experience suggests a significantly higher rate of AKI which may, in part, reflect over cautious fluid replacement in patients with significant insensible fluid losses (eg patients with poor oral intake, fever and GI losses). Although excessive fluid administration may worsen pulmonary injury, it is important to maintain circulating volume and renal perfusion, to reduce the potential for AKI.

Severe COVID pneumonia is associated with abnormal clotting parameters and a higher rate of both venous (25-27%) and arterial (3.7%) thrombosis<sup>11,12</sup>. A significantly elevated D Dimer or prolonged PT/APTT may be associated with greater incidence of VTE<sup>11,12</sup>. The combination of COVID pneumonia and acute pulmonary emboli are likely to worsen hypoxia. Heparin use may reduce 28 day mortality in patients with severe COVID pneumonia and significant sepsis induced coagulopathy<sup>13</sup>.

Recommended indications for admission:

- Hypoxia (saturation under 95%)
- Significant chest X ray findings consistent with COVID
- Tachypnoea

Indicators of poor prognosis

- Older age
- Elevated CRP over 125
- Neutrophil to lymphocyte ratio over 3.1
- Significantly elevated troponin (see local ranges)
- Elevated D Dimer
- Platelets under 100
- Elevated ferritin

Recommendation:

- Stop antiproliferative agents (MMF/azathioprine)
- Consider reducing or stopping CNI
- Consider increasing steroids if currently taking them.
  - There is no evidence for benefit of high dose steroids at this stage
- Oxygen therapy to achieve saturations of 92-96%<sup>16</sup> (unless COPD)
- Regular observations, especially saturations, to monitor for rapid deterioration
- Fluid administration to maintain circulating volume but avoid significant overload
- Consider stopping ACEi/ARB in patients at risk of AKI
- Early VTE prophylaxis with heparin, if no contra-indication
- Consider adjunctive antibiotics if superadded bacterial infection is suspected
- Early discussion of ceilings of care

### **3: Patients who are progressively unwell and require ventilatory support**

Deterioration to requirement for ventilation may occur precipitously and discussion of ceilings of care should start at an early stage. The decision to escalate patients to critical care should be made on an individual basis in line with national guidance and frailty scoring.

It is recommended to start the conversation at an early point in clinical care: <https://www.nice.org.uk/guidance/ng159>

The mortality of all patients requiring critical care support is close to 50%.

There may be a role for the use of methyl prednisolone in patients who develop ARDS<sup>9</sup>

#### Recommendation:

- Stop antiproliferative agents (MMF/azathioprine)
- Dramatically reduce or stop CNI
- Consider high dose steroids in discussion with ITU team
- Ventilatory support in line with local or national guidance
- Adjunctive support or antivirals in line with local practice or clinical trials

#### **4: Additional agents, antivirals and other considerations**

At present, there is no high level evidence of benefit for specific treatments for COVID 19. The use of adjunctive agents or antivirals should be considered in conjunction with local practice or as part of clinical trials.

##### ACEi and ARB

Angiotensin Converting Enzyme 2 (ACE 2) may play a role in coronavirus infection but there is conflicting evidence from basic science studies about the likely effect that modulation of the renin-angiotensin system would have on infection. A number of transplant recipients may be taking ACEi/ARB for their beneficial effect on proteinuria or management of cardiac failure. Therefore it is recommended that the standard advice on continuing or stopping ACEi/ARBs should be adhered to. Temporary discontinuation should be considered in patients who are at risk of Acute Kidney Injury.

The Renal Association guidance can be found here:

<https://renal.org/covid-19/ra-resources-renal-professionals/renal-association-uk-position-statement-patients-novel-corona-virus-infection-use-blood-pressure-medications/>

##### NSAIDs

Reports in the media suggested that NSAIDs, such as Ibuprofen, may worsen the outcome of patients with COVID 19 through the potentiation of ACE 2. Recent statements from international medicine regulators state that there is no evidence to link poor outcomes to NSAIDs and patients who require these agents should continue. As most transplant patients do not use NSAIDs and there are other available antipyretics, we would suggest not discontinuing these agents in patients who require them for their pain or anti-inflammatory properties.

The current UK government guidance on use of NSAIDS is here:

<https://www.gov.uk/government/news/ibuprofen-use-and-covid19coronavirus>

##### Interactions with transplant immunosuppression

Tacrolimus, ciclosporin and sirolimus can interact with some antifungal and antibiotic medications - for example fluconazole and clarithromycin. Where possible use an alternative agent to avoid interaction or reduce dose of immunosuppressant.

Anti-virals e.g. lopinavir/ritonavir (boosted protease inhibitor) have been used in other countries as a possible treatment for COVID 19 and is a treatment arm in RECOVERY Publication version 2, updated 1<sup>st</sup> April 2020 trial. These medicines significantly interact with tacrolimus, ciclosporin, sirolimus and should not be used. The degree of cytochrome P450 enzyme inhibition is profound and can be seen within a few days. Contact transplant specialist for advice.

## References

1. Javier Carbajo-Lozoyaa, Virus Res 165 (2012) 112–117
2. Ma-Lauer Y, Antiviral Res. 2020 Jan;173
3. Ogimi C J Infect Dis. 2017 Jul 15;216(2):203-209
4. Zhang X J Virol. 2008 May;82(9):4420-8.
5. Lee N J Clin Virol. 2004 Dec;31(4):304-9
6. Russell CD Lancet. 2020 Feb 15;395(10223):473-475.
7. Wu Z, JAMA. 2020 Feb 24 epub.
8. Wang D, JAMA. 2020 Feb 7, epub
9. Wu C, JAMA Intern Med. 2020 Mar 13
10. Yichun Cheng,  
<https://www.medrxiv.org/content/10.1101/2020.02.18.20023242v1.article-info>
11. S Chiu, Journal of Thrombosis and Haemostasis <https://doi.org/10.1111/jth.14830>
12. FA Klok, Thrombosis Research <https://doi.org/10.1016/j.thromres.2020.04.013>
13. Tang, Journal of Thrombosis and Haemostasis. [doi:10.1111/JTH.14817](https://doi.org/10.1111/JTH.14817)
14. M Fernandez-Ruiz. AJT. 2020. <https://doi.org/10.1111/ajt.15929>
15. Personal communications – NHS BT and Renal Registry.
16. Clinical guide for the optimal use of Oxygen therapy during the coronavirus pandemic  
<https://www.england.nhs.uk/coronavirus/wp-content/uploads/sites/52/2020/04/C0256-specialty-guide-oxygen-therapy-and-coronavirus-9-april-2020.pdf>

## Appendix 1: Flow chart

